

For a quasi-projective variety S over a field,

$ICH^r(S)$, the intersection Chow group, is defined;

properties (some of which are conjectural) are discussed.

Cf. Barthel, Brasselet, Fieseler, Gabber and Kaup: Relèvement de cycles algébriques et homomorphismes associés en homologie d'intersection, Ann. Math. 141 (1995).

S : variety / \mathbb{C} , $d = \dim S$

$$\begin{array}{ccccc} CHC^r(S) & \longrightarrow & ICH^r(S) & \xrightarrow{\textcircled{1}} & CH_{d-r}(S) \\ \downarrow & & \downarrow & & \downarrow \\ H^{2r}(S) & \longrightarrow & IH^{2r}(S) & \longrightarrow & H_{2(d-r)}^{BM}(S) \end{array}$$

$CH_*(S)$ is Chow group of S ($\otimes \mathbb{Q}$),

$CHC^r(S)$ is Chow cohomology,

① conjecturally exists and surjective.

Thm. [BBFGK]

$$\text{Im} [CH_{d-r}(S) \rightarrow H_{2(d-r)}^{BM}(S)]$$

$$\subset \text{Im} [IH^{2r}(S) \rightarrow H_{2(d-r)}^{BM}(S)] .$$

Let $p : X \rightarrow S$ be a projective map
(with X smooth).

There is a Whitney stratification

$$S = S_0 \supset S_1 \supset \cdots \supset S_\alpha \supset \cdots$$

of S , and resolutions

$$\tilde{X}_\alpha \rightarrow X_\alpha = p^{-1}S_\alpha \text{ such that}$$

$$\tilde{X}_\alpha \rightarrow S_\alpha \text{ is smooth over } S_\alpha - S_{\alpha+1}.$$

$$\begin{array}{ccc}
 & & \tilde{X}_\alpha \\
 & \swarrow \tau_\alpha & \downarrow \\
 X & \supset & X_\alpha \\
 p \downarrow & & \downarrow \\
 S & \supset & S_\alpha
 \end{array}$$

Now take $p : X \rightarrow S$ to be a
resolution of singularities. One has
($d = \dim S$)

$$\mathrm{CH}_{d-r}(\tilde{X}_\alpha) \xrightarrow{\iota_{\alpha*}} \mathrm{CH}^r(X) \xrightarrow{\iota_{\alpha*}} \mathrm{CH}^r(\tilde{X}_\alpha).$$

Each group has a filtration F_S^\bullet (to be explained later).

Define intersection Chow group by:

$$\mathrm{ICH}^r(S) :=$$

$$\frac{\bigcap_{\alpha \geq 1} (\iota_\alpha^*)^{-1} F_S^{2r-d+1} \mathrm{CH}^r(\tilde{X}_\alpha)}{\sum_{\alpha \geq 1} \iota_{\alpha*} F_S^{2r-d+1} \mathrm{CH}_{d-r}(\tilde{X}_\alpha)}$$

Theorem. $\mathrm{ICH}^r(S)$ is well-defined (indep. of choice of stratification and resolution).

There is a map

$$\mathrm{ICH}^r(S) \rightarrow IH^{2r}(S).$$

Bloch, Beilinson, Murre, Shuji Saito
(for case $S = \text{Spec } k$).

Example. X smooth projective
variety.

$CH^r(X)$

$\supset F^1 CH^r(X)$ homologically trivial

$\supset F^2 CH^r(X)$ Kernel of Abel-Jacobi
map ?

Relative canonical filtration. Let X
be smooth, and $p : X \rightarrow S$ be a
projective map.

There is a filtration F_S^\bullet on $CH^r(X)$
satisfying:

$$(1) CH^r(X) = F_S^{-\dim S} CH^r(X).$$

Functorial: for $q : W \rightarrow S$ and

$\Gamma \in \text{CH}_{\dim X+s}(W \times_S X)$, the induced map

$$\Gamma_* : \text{CH}^{r-s}(W) \rightarrow \text{CH}^r(X)$$

respects F_S^\bullet .

(2) If the induced map $[\Gamma] :$

$${}^p\mathcal{H}^{2r+2s-\nu} Rq_* \mathbb{Q}_W \rightarrow {}^p\mathcal{H}^{2r-\nu} Rp_* \mathbb{Q}_X$$

is zero, then Γ_* sends F_S^ν to $F_S^{\nu+1}$.

(3) The filtration is the smallest with properties (1) and (2).

Proposition. Under Conjectures,
 $F_S^\nu \text{CH}^r(X) = 0$ for $\nu \gg 0$.

Theorem 1 (Under Conjectures)
There is a natural surjective map
 $\text{ICH}^r(S) \rightarrow \text{CH}^r(S)$.


Theorem 2. (Without Conjectures)

$$\text{Im} [\text{CH}_{d-r}(S) \rightarrow H_{2(d-r)}^{\text{BM}}(S)]$$

$$\subset \text{Im} [\text{IH}^{2r}(S) \rightarrow H_{2(d-r)}^{\text{BM}}(S)]$$

Proof of Thm 1. in special case.

Assume :



$$\dim Y = d' , \quad \dim q = e$$

$$\dim \Sigma = d' - e$$

Must show: For $\forall a \in CH_{d+1}(S)$

$\exists b \in CH^r(X)$ s.t.

(i) $(z^*)^{-1} b \in F_S^{2r-d+1} CH^r(\tilde{Y})$,

(ii) $p_*(b) = a$.

To prove, take any b satisfying (ii), and modify by $z_*(c)$ using the following three lemmas.

Under conjectures,

① In the sequence

$$\mathrm{Gr}_{F_S}^{\nu} \mathrm{CH}_{d-r}^r(\tilde{y}) \xrightarrow{L^*} \mathrm{Gr}_{F_S}^{\nu} \mathrm{CH}^r(X) \xrightarrow{L^*} \mathrm{Gr}_{F_S}^{\nu} \mathrm{CH}^r(\tilde{y})$$

L^* is injective for $\nu \neq 2r-d$

L_* is surjective for $\nu \neq 2r-d$.

$$\textcircled{2} \quad \mathrm{CH}^r(\tilde{y}) = F_S^{2r-d'-e} \mathrm{CH}^r(\tilde{y})$$

[perverse degree of $Rq_* \mathbb{Q}_y$
is in $[d'-e, d'+e]$]

$$\textcircled{3} \quad F_S^{2r-d'-e} \mathrm{CH}_{d-r}(\Sigma) = 0$$

[perv. degree of D_{Σ} is $z - \dim \Sigma$]

Conjectures

Grothendieck's Standard conjecture
(\Rightarrow semi-simplicity of the category of pure homological motives).

Bloch-Beilinson-Murre: Existence of Chow-Künneth decomposition (with properties...)

($\Rightarrow h(X) = \bigoplus h^i(X)$ in the category of Chow motives.)

Beilinson-Soulé: vanishing of motivic cohomology with negative degree.

Topological theory:

$D_c^b(S)$: derived category of sheaves with cohomology constructible sheaves;

For a map $f : S \rightarrow S'$, f^* , f_* , $f^!$, $f_!$;

Poincaré-Verdier duality formulas;

For $p : X \rightarrow S$,

$$H^i(X, \mathbb{Z}) = \text{Hom}(\mathbb{Z}_S, R p_* \mathbb{Z}_X[i]).$$

perverse t -structure. In particular, perverse cohomology functors

$${}^p\mathcal{H}^\nu : D_c^b(S) \rightarrow \text{Perv}(S).$$

Motivic theory:

$\mathcal{D}(S)$: triangulated category of motives over S :

For a map f , f^* , f_* , $f^!$, $f_!$;

For $p : X \rightarrow S$, $H_{\mathcal{M}}^i(X, \mathbb{Z}(r)) = \text{Hom}(\mathbb{Z}_S(0), R p_* \mathbb{Z}_X(r)[i])$.

Poincaré-Verdier duality formulas;

perverse t -structure.

Realization functor

$$\rho : \mathcal{D}(S) \rightarrow D_c^b(S).$$

Theorem. (assume $\text{ch} = 0$ for simplicity) There is a triangulated category $\mathcal{D}(S)$ (called the category of mixed motives over S) with properties:

(1) There is a functor $h :$
 $(\text{Quasi-Projective } /S)^{opp} \rightarrow \mathcal{D}(S)$

There are Tate objects $\mathbb{Z}_S(r)$.

(2) Natural isomorphism

$$\text{Hom}(\mathbb{Z}_S(0), h(X/S) \otimes \mathbb{Z}(r)[2r-n]) = \text{CH}^r(X, n) .$$

In particular,

$$\text{Hom}(\mathbb{Z}_S(0), h(X/S) \otimes \mathbb{Z}_S(r)[2r]) = \text{CH}^r(X) .$$

(4) There are functors $\otimes, f^*, f_*, f^!, f_!$ among the categories $\mathcal{D}(S)$,

satisfying the correct properties (such as Verdier duality).

(5) ($k \subset \mathbb{C}$) There is the realization functor

$$\rho : \mathcal{D}(S) \rightarrow D_c^b(S(\mathbb{C}))$$

such that $\rho \circ h$ is the cohomology functor for varieties.

From now, write $\mathcal{D}(S)$ for $\mathcal{D}(S)_{\mathbb{Q}}$.

Theorem. (Under the conjectures of Grothendieck, Bloch-Beilinson-Murre, and Beilinson-Soulé)

(1) There is a Whitney stratification $\{S_{\alpha}\}$ of S , local systems \mathcal{V}_{α}^i on $S_{\alpha} - S_{\alpha+1}$, and a non-canonical direct sum decomposition

$$h(X/S) = \bigoplus_{i,\alpha} h_{\alpha}^i(X/S) \text{ in } \mathcal{D}(S)$$

such that $\rho(h_\alpha^i(X/S)) \cong IC_{S_\alpha}(\mathcal{V}_\alpha^i)[-i + \dim S_\alpha]$.

(Work with Corti)

(2) There is a t -structure on $\mathcal{D}(S)_\mathbb{Q}$ such that ρ is compatible with it and the perverse t -structure on $D_c^b(S)$.

(As a consequence, there is an abelian subcategory $MM(S)$, and functors

$${}^p\mathcal{H}^\nu : \mathcal{D}(S) \rightarrow MM(S).$$

The category $MM(S)$ is abelian, and the induced functor

$MM(S) \rightarrow Perv(S)$ is exact and faithful.